


AMENDMENTS TO THE CLAIMS

1. (Original) A fungicidal mixture comprising, as active components,


- 1) the triazolopyrimidine derivative of the formula I



I

and

- 2) epoxiconazole of the formula II,



II

in a synergistically effective amount.

2. (Original) A fungicidal mixture comprising the compound of the formula I and the compound of the formula II in a weight ratio of from 100:1 to 1:100.
3. (Currently amended) A fungicidal composition comprising a liquid or solid carrier and a mixture as claimed in claim 1 or 2.
4. (Original) A method for controlling harmful fungi from the class of the *Oomycetes*, which comprises treating the fungi, their habitat or the seed, the soil or the plants to be protected against fungal attack with an effective amount of the compound I and the compound II as set forth in claim 1.

5. (Currently amended) A method as claimed in claim 4, wherein the compounds I and II as ~~set forth in claim 1~~ are applied simultaneously, that is jointly or separately, or in succession.
6. (Currently amended) A method as claimed in claim 4, wherein the mixture ~~as claimed in claim 1 or 2~~ of the compound I and the compound II is applied to the soil or the plants to be protected against fungal attack in an amount of from 5 g/ha to 2000 g/ha.
7. (Currently amended) A method as claimed in claim 4 ~~or 5~~, wherein the mixture ~~as claimed in claim 1 or 2~~ of the compound I and the compound II is applied in an amount of from 0.001 to 1 g/kg of seed.
8. (Currently amended) A method as claimed in ~~any of claims 4 to 7~~ claim 4, wherein the harmful fungus *Phytophthora infestans* is controlled.
9. (Currently amended) Seed comprising the mixture as claimed in claim 1 ~~or 2~~ in an amount of from 0.001 to 1 g/kg.
10. (Original) The use of the compound I and the compound II as set forth in claim 1 for preparing a composition suitable for controlling *Oomycetes*.
11. (New) A fungicidal composition comprising a liquid or solid carrier and a mixture as claimed in claim 2.
12. (New) The method as claimed in claim 6, wherein the mixture comprises the compound of the formula I and the compound of the formula II in a weight ratio of from 100:1 to 1:100.
13. (New) The method as claimed in claim 7, wherein the mixture comprises the compound of the formula I and the compound of the formula II in a weight ratio of from 100:1 to 1:100.
14. (New) A method as claimed in claim 5, wherein the harmful fungus *Phytophthora infestans* is controlled.

15. (New) A method as claimed in claim 6, wherein the harmful fungus *Phytophthora infestans* is controlled.
16. (New) A method as claimed in claim 7, wherein the harmful fungus *Phytophthora infestans* is controlled.
17. (New) Seed comprising the mixture as claimed in claim 2 in an amount of from 0.001 to 1 g/kg.
18. (New) A method as claimed in claim 5, wherein the mixture of the compound I and the compound II is applied in an amount of from 0.001 to 1 g/kg of seed.
19. (New) The method as claimed in claim 18, wherein the mixture comprises the compound of the formula I and the compound of the formula II in a weight ratio of from 100:1 to 1:100.